arXiv:1205.0759v1 [math.CV] 3 May 2012

On H^∞ on the complement of $C^{1+\alpha}$ curves

J.M. Enríquez-Salamanca[b,**], María J. González[a,1,*]

[a] Department of Mathematics, University of Cádiz, Puerto Real, 11510, Spain
[b] Department of Mathematics, University of Cádiz, Cádiz, 11002, Spain

---

Abstract

Let ρ be a quasiconformal mapping on the plane with complex dilatation μ. We show that if μ satisfies a certain Carleson measure condition, then one can transfer H^∞ on the upper half plane onto the corresponding space in the complement of the quasicircle $\Gamma = \rho(\mathbb{R})$, and that this condition on μ characterizes $C^{1+\alpha}$ curves.

Keywords: Cauchy integral; quasiconformal mapping; Carleson measures; smooth curves

---

Introduction

Let f be a quasiconformal self-mapping of the plane with complex dilatation μ. Thus f is a homeomorphism with locally integrable distributional derivatives verifying that $\bar{\partial} f - \mu \partial f = 0$, $\mu \in L^\infty(\mathbb{C})$ and $\|\mu\|_\infty < 1$. We shall also assume throughout this paper that μ has compact support.

The images of the real line (or the unit circle $\partial\mathbb{D}$) under quasiconformal mappings of the plane are called quasicircles. In general, they are not rectifiable and they do not satisfy any regularity conditions such as local absolute continuity or differentiability a.e., even when $\|\mu\|_\infty$ is small. Then, understanding the properties of the geometry of quasicircles in terms of the complex dilatation becomes one of the main objectives of the quasiconformal

---

[*] Corresponding author

[**] Principal corresponding author

Email addresses: `enrique.desalamanca@uca.es` (J.M. Enríquez-Salamanca), `majose.gonzalez@uca.es` (María J. González)

[1] This author is supported by Grant MTM 2001-24606.

analysis and, also, of this article.

One can get some regularity by imposing some stronger smallness condition on the dilatation μ. If μ were zero in a neighborhood of $\partial\mathbb{D}$, then the map would be smooth on $\partial\mathbb{D}$. So, if μ decays to zero in some sense as it approaches the unit circle then we should be able to get some good behaviour of the mapping on $\partial\mathbb{D}$.

One of the first results along these lines is due to Carleson [6]. He showed that if f is a quasiconformal self mapping of $\mathbb{C}\backslash\mathbb{D}$ such that

$$\int_0^1 \frac{K(t)^2}{t}dt < \infty,$$

where $K(t) = \sup\{|\mu(z)| : 1 < |z| < 1+t\}$, then f is absolutely continuous on the circle and $f' \in L^2_{loc}$. Becker [3] extended this result to the case where f represents a conformal mapping of $\mathbb{D}$ that extends quasiconformally to the whole plane.

In the same context, Dynkin [7] proved the following stronger result. Let

$$Q(\mu)(z) = \left(\int_{1<|\zeta|<2} \frac{|\mu(\zeta)|^2}{|\zeta - z|^2} d\xi d\eta\right)^{1/2}.$$

If there exists a constant $a > 0$ so that $e^{aQ(\mu)^2} \in L^1(\partial\mathbb{D})$, then the curve $\Gamma = f(\partial\mathbb{D})$ is rectifiable.

A particular type of rectifiable quasicircle are the chord-arc or Laurentiev curves. A Jordan curve, Γ, is M chord-arc if satisfies $\Lambda(\Gamma(z_1, z_2)) \leq M|z_1 - z_2|$, where $\Lambda(\Gamma(z_1, z_2))$ denotes the length of the shortest arc of the curve Γ between $z_1, z_2 \in \Gamma$.

A condition for chord-arc curves with small constant was given by Astala-Zinsmeister [2] , and requires the measure

$$\frac{|\mu(z)|^2}{|z| - 1} dxdy$$

to be a Carleson measure in $\mathbb{C}\backslash\mathbb{D}$ with small norm. The proof relies on estimates for the Schwarzian derivative of f in $\mathbb{D}$, showing that $\log f' \in$ BMOA with small BMO constant. On the other hand Semmes [13] and MacManus [10] showed that the same result holds with no assumption on f being conformal on $\mathbb{D}$. Semmes' proof is based on estimates for a certain perturbed Cauchy integral operator while the strategy in [10] is to estimate Haar type coefficients for $\log f'$.

For arbitrary constants, the result is no longer true [5]. In fact, if no restriction on the Carleson norm of $|\mu(z)|^2/(|z|-1)$ is imposed, the quasicircle Γ might not be even rectifiable.

Related to these results is the following one due to Pommerenke [11]. A curve Γ is an asymptotically smooth curve, that is

$$\frac{\Lambda(\Gamma(z_1,z_2))}{|z_1-z_1|} \to 1, \quad \text{as} \quad |z_1-z_2| \to 0,$$

if and only if $\log f' \in \mathrm{VMOA}$.

This paper presents a new characterization of smooth curves in terms of μ. In particular, we are interested in $C^{1+\alpha}$ curves. We prove the following result:

Theorem 1. *Let f denote a conformal map of $\mathbb{D}$ onto the inner domain of a Jordan curve Γ. Then Γ is a $C^{1+\alpha}$ curve if and only if f extends to a global quasiconformal map whose dilatation μ satisfies that $|\mu(z)|^2/(|z|-1)^{1+\varepsilon}$ is a Carleson measure relative to $\partial\mathbb{D}$, where $\varepsilon = \varepsilon(\|\mu\|_\infty)$ and $\alpha = \alpha(\varepsilon, \|\mu\|_\infty)$.*

The equivalent result holds if we consider unbounded $C^{1+\alpha}$ curves. In this case μ satisfies that $|\mu(z)|^2/|y|^{1+\varepsilon}$ is a Carleson measure relative to $\mathbb{R}$, where $y = \mathrm{Im}(z)$.

In the second part of the paper we consider quasiconformal mappings of the plane $\rho : \mathbb{C} \to \mathbb{C}$ whose complex dilatation μ satisfies that for some $\varepsilon > 0$, $|\mu|^2/|y|^{1+\varepsilon}$ is a Carleson measure relative to $\mathbb{R}$. We will show that under this condition on μ we can transfer H^∞ on the half plane $\mathbb{R}^2_+$ onto the corresponding space in the complement of the quasicircle $\Gamma = \rho(\mathbb{R})$. More precisely, we prove the following result:

Theorem 2. *Let ρ be a quasiconformal map of the plane onto itself whose complex dilatation μ satisfies that for some $\varepsilon > 0$, $|\mu|^2/|y|^{1+\varepsilon}$ is a Carleson* *measure relative to* $\mathbb{R}$. *Let* Ω_+ *and* Ω_- *denote the two regions bounded by the quasicircle* $\Gamma = \rho(\mathbb{R})$. *Then* $G \in H^\infty(\Omega_\pm)$ *if and only if* $F \in H^\infty(\mathbb{R}^2_\pm)$ *respectively, where* $F|_{\mathbb{R}} = G \circ \rho|_{\mathbb{R}}$.

The proof is based on an argument by Semmes [12], where the idea is to transform a $\bar\partial$ problem relative to Γ into a $\bar\partial - \mu\partial$ problem relative to $\mathbb{R}$ via a change of variables.

We will apply the previous results to study the relation between the Cauchy integral $C_\Gamma(g)$ of a function $g \in L^\infty(\Gamma)$, that is

$$C_\Gamma(g)(z) = \frac{1}{2\pi i}\int_\Gamma \frac{g(\omega)}{\omega - z}d\omega, \quad z \notin \Gamma,$$

and the Cauchy integral of its pullback on $\mathbb{R}$. So, as an immediate consequence of Theorem 1 and Theorem 2 we obtain the following corollary.

Corollary. *Let* Γ *be an unbounded* $C^{1+\alpha}$ *curve analytic at* ∞ *and let* $g \in L^\infty(\Gamma)$. *Then the Cauchy integral* $C_\Gamma(g) \in L^\infty(\mathbb{C})$ *if and only if* $C_{\mathbb{R}}(f) \in L^\infty(\mathbb{C})$ *where* f *denotes the pullback of* g *under the conformal mapping that sends* $\mathbb{R}$ *onto* Γ.

The paper is structured as follows: In section 1, we review some definitions and basic facts, in particular the analytic characterization of $C^{1+\alpha}$ curves. The proof of Theorem 1 is presented in section 2 whereas section 3 is devoted to Theorem 2.

1. Preliminaries.

Let us denote complex variables by $z = x + iy$ and $\zeta = \xi + i\eta$. We shall use the following notation throughout this article: $\mathbb{D} = \{z : |z| < 1\}$, $\mathbb{T} = \partial\mathbb{D}$, $B_x(r)$ denotes the ball centered at x and radius r, $|I|$ represents the length of any arc $I \subset \partial\mathbb{D}$, $\Gamma(z_1, z_2)$ is the shortest arc of the curve Γ between $z_1, z_2 \in \Gamma$, and $\Lambda(\Gamma)$ the length of the curve Γ. Also, we shall write $\bar\partial = \partial/\partial\bar z = 1/2(\partial_x + i\partial_y)$ and $\partial = \partial/\partial z = 1/2(\partial_x - i\partial_y)$.

A positive measure λ on $\mathbb{C}$ is called a Carleson measure relative to a given chord-arc curve Γ if there exists a constant $C > 0$ such that $\lambda(B_z(R)) \leq C\ R$ for all $z \in \Gamma$ and $R > 0$. The smallest such C is the norm of λ, $\|\lambda\|_C$.

A function $f \in L^1(\mathbb{T})$ belongs to the space $\mathrm{BMO}(\mathbb{T})$ if there exists $A > 0$ so that

$$\sup_I \frac{1}{\Lambda(I)} \int_I |f(\zeta) - a_I||d\zeta| \leq A, \text{ with } a_I = \frac{1}{\Lambda(I)} \int_I f(\zeta)|d\zeta|$$

and where the supremum is taken over all arcs $I \subseteq \mathbb{T}$. The least possible A in this inequality is called the BMO norm of f, $\|f\|_*$.

The space VMO is defined as follows:

$$\mathrm{VMO}(\mathbb{T}) = \{f \in \mathrm{BMO}(\mathbb{T}) : \lim_{|I| \to 0} \frac{1}{\Lambda(I)} \int_I |f(\zeta) - a_I||d\zeta| = 0\}.$$

We say that $f \in$ BMOA or $f \in$ VMOA if $f \in H^1(\mathbb{D})$ and if the boundary values of f on $\mathbb{T}$ belong to $\mathrm{BMO}(\mathbb{T})$ or $\mathrm{VMO}(\mathbb{T})$ respectively. Recall that BMOA is contained in the Bloch space

$$\mathcal{B} = \{f \text{ analytic in } \mathbb{D} : \|f\|_{\mathcal{B}} = \sup(1 - |z|^2)|f'(z)| < \infty\}$$

and VMOA is contained in $\mathcal{B}_0 = \{f \in \mathcal{B} : (1 - |z|^2)|f'(z)| \to 0 \text{ as } |z| \to 1-0\}$.

The notion of Carleson measures is closely related to BMO functions. Let F be an analytic function on $\mathbb{R}^2_+$ and $f = F|_{\mathbb{R}}$. Then $f \in \mathrm{BMO}(\mathbb{R})$ if and only if $|F'(z)|^2 y\, dxdy$ is a Carleson measure with respect to $\mathbb{R}$ [8]. Analogous definitions and results extend to analytic functions defined on the complement of chord-arc curves.

We now introduce the curves which are the main object of study in this paper. A Jordan curve Γ is said to be of class C^n ($n = 1, 2, \ldots$) if it has a parametrization $\varphi(\tau) = f(e^{i\tau}), 0 \leq \tau \leq 2\pi$, that is n times continuously differentiable and satisfies that $\varphi'(\tau) \neq 0$, $\forall \tau$. Furthermore, it is of class $C^{n,\alpha}$, for $0 < \alpha < 1$, if

$$|\varphi^{(n)}(\tau_1) - \varphi^{(n)}(\tau_2)| \leq C|\tau_1 - \tau_1|^\alpha \tag{1}$$

It is well known that for $0 < \alpha < 1$ we can consider the parametrization of the curve given by the conformal mapping f that sends $\mathbb{D}$ onto the inner domain bounded by Γ (Kellogg-Warschawski Theorem). In this case, (1) is equivalent to

$$|f^{n+1}(z)| \leq C(1 - |z|)^{\alpha - 1}, \text{ for all } z \in \mathbb{D}. \tag{2}$$

2. Proof of Theorem 1.

The main idea is based on an estimate of the logarithmic derivative developed by Dynkin [7]. Accordingly, if f is a conformal mapping in the unit disc with a k-quasiconformal extension to the whole plane, that is $\|\mu\|_\infty \leq k < 1$, then for all $z \in \mathbb{D}$

$$(1-|z|)\left|\frac{f''(z)}{f'(z)}\right| \leq C(1-|z|)^{1-k}\left[1+\int_{1-|z|}^{1}\frac{w(z,t)}{t^{2-k}}dt\right], \tag{3}$$

where

$$\omega(z,t)=\left(\frac{1}{\pi t^2}\int_{|\zeta-z|<t}|\mu(\zeta)|^2 d\xi d\eta\right)^{1/2}. \tag{4}$$

Using this estimate, he also proved the following result:

Theorem 3. *If the integral*

$$\int_0^1 \frac{\omega(z,t)}{t}dt < \infty$$

converges uniformly in $z \in \partial\mathbb{D}$*, then* $\log f'$ *(and thereby* f' *and* $1/f'$*) is continuous in the closed disc.*

Theorem 1. *Let* f *denote a conformal map of* $\mathbb{D}$ *onto the inner domain of a Jordan curve* Γ*. Then* Γ *is a* $C^{1+\alpha}$ *curve if and only if* f *extends to a global quasiconformal map whose dilatation* μ *satisfies that* $|\mu(z)|^2/(|z|-1)^{1+\varepsilon}$ *is a Carleson measure relative to* $\partial\mathbb{D}$*, where* $\varepsilon=\varepsilon(\|\mu\|_\infty)$ *and* $\alpha=\alpha(\varepsilon,\|\mu\|_\infty)$*.*

Proof. Let us assume first that there exists a quasiconformal extension of f so that for some $\varepsilon>0$ $\nu(z)=|\mu(z)|^2/(|z|-1)^{1+\varepsilon}$ is a Carleson measure on the plane, $\|\mu\|_\infty \leq k < 1$. Let $z\in\mathbb{D}$, then for any $t>1-|z|$ we obtain from (4) that:

$$\begin{aligned}\omega(z,t)^2 &\leq \frac{1}{\pi t^2}\left(\int_{B_z(t)}\frac{|\mu(\zeta)|^2}{(|\zeta|-1)^{1+\varepsilon}}d\xi d\eta\right)^{1/2}\left(\int_{B_z(t)}|\mu(\zeta)|^2(|\zeta|-1)^{1+\varepsilon}d\xi d\eta\right)^{1/2}\\ &\leq Ct^{\varepsilon/2}.\end{aligned} \tag{5}$$

where $C=C(\|\mu\|_\infty,\|\nu\|_C)$. Therefore, if $z\leq \min(\varepsilon/4;1-k)$, by (3)

$$\left|\frac{f''(z)}{f'(z)}\right| \leq C(1-|z|)^{-k}\left[1+C\int_{1-|z|}^{1} t^{\varepsilon/4+k-2}dt\right] \leq C(1-|z|)^{\alpha-1}$$

and $|f''(z)| \leq C|f'(z)|(1-|z|)^{\alpha-1}$, where now $C = C(\varepsilon, \|\mu\|_\infty, \|\nu\|_C)$.

We only need to prove that $|f'(z)|$ is bounded on $\mathbb{T}$. Since

$$\int_0^1 \frac{\omega(z,t)}{t}dt \leq C\int_0^1 \frac{t^{\varepsilon/4}}{t}dt < \infty,$$

and by Theorem 3, $|f'|$ is bounded in the closed disc. Therefore, $|f''(z)| \leq C(1-|z|)^{\alpha-1}$ which implies by (2) that Γ is $C^{1+\alpha}$.

To prove the second part of the theorem, let us consider the following quasiconformal extension of the Riemann mapping f:

$$f(z) = f(\frac{1}{\bar{z}}) + f'(\frac{1}{\bar{z}})(z-\frac{1}{\bar{z}}), \quad \text{for} \quad |z|>1,$$

where

$$f_z = f'(\frac{1}{\bar{z}}) \quad \text{and} \quad f_{\bar{z}} = -\frac{1}{\bar{z}^2}(z-\frac{1}{\bar{z}})f''(\frac{1}{\bar{z}}).$$

Note that if $\beta(z)$ denotes the logarithmic derivative, that is $\beta(z) = (1-|z|)|f''(z)/f'(z)|$, $z\in\mathbb{D}$, then $|\mu(1/\bar{z})| \asymp \beta(z)$.

In general, this mapping is not homeomorphic, but Becker and Pommerenke [4] proved that this extension was indeed a quasiconformal extension of f to a neighborhood of $\mathbb{T}$ if $\overline{\lim}_{|z|\to 1-0}\beta(z) < 1$.

If Γ is $C^{1+\alpha}$, it is assymptotically smooth, so by Pommerenke's result $\log f' \in$ VMOA. Since VMOA $\subset \mathcal{B}_0$, we get that $\beta(z)\to 0$ as $|z|\to 1-0$ and the aforementioned extension of f defines a well defined quasiconformal extension to a neighborhood of $\mathbb{T}$.

As $|\mu(1/\bar{z})| \asymp \beta(z)$ and Γ is a $C^{1+\alpha}$ curve, for $|z|>1$

$$|\mu(z)| \leq C\beta(\frac{1}{\bar{z}}) \leq C(|z|-1)^{\alpha}.$$

A straightforward computation shows that, for $\varepsilon \leq 2\alpha$, $z \in \partial\mathbb{D}$ and $R > 0$

$$\int_{B_z(R)} \frac{|\mu(\zeta)|^2}{(|\zeta|-1)^{1+\varepsilon}} d\xi d\eta \leq C \int_{B_z(R)} \frac{(|\zeta|-1)^{2\alpha}}{(|\zeta|-1)^{1+\varepsilon}} d\xi d\eta \leq CR.$$

$\square$

3. Proof of Theorem 2.

Before proceeding to the proof we need to mention a result on quasiconformal mappings preserving Carleson measures. We say that $F : \mathbb{R}^2_+ \to \mathbb{R}^2_+$ preserves Carleson measures if given any Carleson measure μ in $\mathbb{R}^2_+$, the measure ν defined in $\mathbb{R}^2_+$ as

$$\nu(E) = \int_{F^{-1}(E)} a_F(z) d\mu(z)$$

is a Carleson measure, where

$$a_F(z) = \frac{1}{|B_z|} \int\int_{B_z} J_F(\zeta)^{1/2} d\zeta \wedge \bar{\zeta},$$

$B_z = B_z(1/2\ y)$ and J_F is the Jacobian of F. The function $a_F(z)$ is somehow the quasiconformal substitute of $|F'|$ in Koebe distortion theorem. In fact $\operatorname{Im} F(z) \simeq a_F(z) y$ [1].

If we consider a quasiconformal map ρ from $\mathbb{R}^2_+$ onto itself, it is well known that ρ preserves Carleson measures if $|\rho'| \in A_\infty$ [2], [9]. We will need the analogous result for a quasiconformal map ρ from $\mathbb{R}^2_+$ onto a domain bounded by a chord-arc curve.

Lemma 1. *Let $\rho : \mathbb{R}^2_+ \to \Omega$ a quasiconformal map and let $\partial\Omega$ be a chord-arc curve. If $|\rho'| \in A_\infty$ then ρ preserves Carleson measures.*

The proof follows from the previous case, that is $\Omega = \mathbb{R}^2_+$, since Ω is the image of the upper half plane under a global bilipschitz map.

To prove Theorem 2 we will follow Semmes approach in [12]. Let Γ be locally rectifiable quasicircle in the plane. Given a function g defined on Γ, consider its Cauchy integral

$$C_\Gamma(g)(z) = \frac{1}{2\pi i}\int_\Gamma \frac{g(\omega)}{\omega - z} d\omega, \quad z \notin \Gamma.$$

We define the jump of $G = C_\Gamma(g)$ across Γ at the point z as $g_+ - g_-$, where g_+ and g_- denote the boundary values of G. As the classical Plemelj formula states,

$$g_\pm(z) = \pm\frac{1}{2}g(z) + \frac{1}{2\pi i}P.V.\int_\Gamma \frac{g(\omega)}{\omega - z} d\omega, \quad z \in \Gamma.$$

Hence $g_+(z) - g_-(z) = g(z)$. Also, G is holomorphic off Γ, so that $\bar\partial G = 0$ on $\mathbb{C}\backslash\Gamma$.

We can reexpress these two conditions by saying that, in the distributional sense, $\bar\partial G = g\; dz_\Gamma$ on $\mathbb{C}$, that is

$$\int_\Gamma g\; dz_\Gamma = \int_\Omega (\bar\partial G)\; d\bar z \wedge dz,$$

where dz_Γ is the usual measure on Γ and $\partial\Omega = \Gamma$.

We can also say that G is determined by the equation $\bar\partial G = 0$ on $\mathbb{C}\backslash\Gamma$ and the condition jump$(G) = g$ on Γ, as if $\tilde G$ were another function with the same properties, then $G - \tilde G$ would be entire and a mild condition at ∞ would force it to be 0.

Let $\tilde G = G \circ \rho$ on $\mathbb{C}\backslash\mathbb{R}$ and $f = g \circ \rho$ on $\mathbb{R}$, where $\rho : \mathbb{C} \to \mathbb{C}$ is a quasiconformal map that takes $\mathbb{R}$ into Γ. Then, $\bar\partial G = 0$ off Γ transforms into $(\bar\partial - \mu\partial)\tilde G = 0$ off $\mathbb{R}$ with jump$(\tilde G) = f$ across $\mathbb{R}$, where $\mu = \mu_\rho$. Again, in the distributional sense, we can say that $(\bar\partial - \mu\partial)\tilde G = fdx$.

In order to prove the following result it is convenient to change the problem a bit more. Let $F = C_\mathbb{R}(f)$ the Cauchy integral on $\mathbb{R}$ of f. Thus, F is holomorphic off $\mathbb{R}$ and its jump across $\mathbb{R}$ is given by f, i.e., $\bar\partial F = fdx$.

Let us now define $H = \tilde G - F$ on $\mathbb{C}\backslash\mathbb{R}$. Then, H has no jump across $\mathbb{R}$ and $\bar\partial H = \mu\partial\tilde G$. As H has no jump, we can consider that the previous equation holds on all of $\mathbb{C}$ in the sense of distributions (that there is no boundary piece).

Theorem 2. *Let ρ be a quasiconformal map of the plane onto itself whose complex dilatation μ satisfies that for some $\varepsilon > 0$, $|\mu|^2/|y|^{1+\varepsilon}$ is a Carleson measure relative to $\mathbb{R}$. Let Ω_+ and Ω_- denote the two regions bounded by the quasicircle $\Gamma = \rho(\mathbb{R})$. Then $G \in H^\infty(\Omega_\pm)$ if and only if $F \in H^\infty(\mathbb{R}^2_\pm)$ respectively, where $F|_{\mathbb{R}} = G \circ \rho|_{\mathbb{R}}$.*

Proof. Let $G \in H^\infty(\Omega_+)$ (the same argument would hold for $H^\infty(\Omega_-)$). Then $G = C_\Gamma(g)$ where $g = G|_\Gamma \in L^\infty(\Gamma)$. To show that $F \in \mathbb{H}^\infty(\mathbb{R}^2_+)$ we need to prove that $F|_{\mathbb{R}} \in L^\infty(\mathbb{R})$. Using the notation above, this is equivalent to prove that $H|_{\mathbb{R}} \in L^\infty(\mathbb{R})$. Since $\bar\partial H = \mu\partial\tilde{G}$,

$$H(a) = \frac{1}{\pi i}\int_{\mathbb{C}} \frac{\bar\partial H(z)}{z-a}dxdy = \frac{1}{\pi i}\int_{\mathbb{C}} \frac{\mu(z)\partial\tilde{G}(z)}{z-a}dxdy \text{ for } a \in \mathbb{R}. \tag{6}$$

For $a \in \mathbb{R}$ and k an integer, let us denote $B_k = B_a(2^{-k})$. Then

$$\begin{aligned} |H(a)| &\lesssim \sum_k 2^{k+1}\int_{B_k\setminus B_{k+1}} |\mu(z)||\partial\tilde{G}(z)|dxdy \\ &\lesssim \sum_k 2^{k+1}\left(\int_{B_k}\frac{|\mu(z)|^2}{|y|}dxdy\right)^{1/2}\left(\int_{B_k}|\partial\tilde{G}(z)|^2|y|dxdy\right)^{1/2}. \end{aligned} \tag{7}$$

Since $\nu(z) = |\mu(z)|^2/|y|^{1+\varepsilon}$ is a Carleson measure,

$$\int_{B_k}\frac{|\mu(z)|^2}{|y|}dxdy = \int_{B_k}\frac{|\mu(z)|^2}{|y|^{1+\varepsilon}}|y|^\varepsilon dxdy \le \|\nu\|_C 2^{-k(1+\varepsilon)}. \tag{8}$$

The above inequality also shows that $\tau = |\mu|^2/|y|$ is a Carleson measure with Carleson norm $\|\tau\|_C \to 0$ as $k \to \infty$. In this case, the quasicircle $\Gamma = \rho(\mathbb{R})$ is actually a chord-arc curve with small constant and $\rho|_{\mathbb{R}}$ satisfies that $|\rho'| \in A_\infty$ [10]. On the other hand, since $G \in H^\infty(\Omega_+)$ then $|G'(z)|^2\delta_\Gamma(z)dxdy$ is a Carleson measure relative to Γ with norm $\le C\|g\|_*^2$.

By Lemma 1, $\lambda = |\partial\tilde{G}|^2 y\, dxdy$ will be a Carleson measure relative to $\mathbb{R}$ and

$$\int_{B_k}|\partial\tilde{G}(z)|^2 ydxdy \le \|\lambda\|_C\, 2^{-k}. \tag{9}$$

Then by (7), (8) and (9) $|H(a)| \le C(\|\mu\|_C, \|\lambda\|_C)\sum_k \left(2^{-\varepsilon/2}\right)^k < \infty$ as we wanted to prove.

Conversely, if F were bounded, the same argument would show that $\tilde{G}$ is bounded on $\mathbb{R}$ and then, that $G \in H^\infty(\Omega_+)$. $\square$

To conclude this paper we would like to propose the following problem:

Problem. *Find conditions on μ so that Theorem 2 holds.*

References

[1] K. Astala, F.W. Gehring, Quasiconformal analogues of theorems of Koebe and Hardy-Littlewood, Mich. Math. J. 32 (1985) 99-107.

[2] K. Astala, M. Zinsmeister, Teichmüller spaces and BMOA, Math. Ann. 289 (1991) 613-625.

[3] J. Becker, On asymptotically conformal extension of univalent functions, Complex Variables. 9 (1987) 109-120.

[4] J. Becker, Ch. Pommerenke, ber die quasikonforme Fortsetzung schlichte Funktionen, Math. A. 161 (1978) 69-80.

[5] C. Bishop, P.W. Jones, Harmonic measure, L^2 estimates and the Schwarzian derivative, J. Analyse Math. 62 (1994) 77-113.

[6] L. Carleson, On mappings, conformal at the boundary, J. Analyse Math. 19 (1967) 1-13.

[7] E. Dynkin, Estimates for asymptotically conformal mappings, Annales Academiae Scientiarum Fennicae. 22 (1997) 275-304.

[8] J. Garnett, Bounded Analytic Functions, Academic Press, New York, 1981.

[9] M.J. González, A. Nicolau, Quasiconformal mappings preserving interpolating sequences, Ann. Acad. Sci. Fenn. 23 (1998) 283-290.

[10] P. MacManus, Quasiconformal mappings and Ahlfors-David curves, Trans. Amer. Math. Soc. 343 (1994) 853-881.

[11] Ch. Pommerenke, Boundary Behaviour of Conformal Maps, Springer-Verlag, Berlin, 1973.

[12] S. Semmes, Estimates for $(\bar{\partial} - \nu\partial)^{-1}$ and Calderon's theorem on the Cauchy integral, Trans. Amer. Math. Soc. 306 (1988) 191-232.

[13] S. Semmes, Quasiconformal mappings and chord-arc curves, Trans. Amer. Math. Soc. 306 (1988) 233-263.